




AENSI Journals

Advances in Environmental Biology

ISSN-1995-0756 EISSN-1998-1066

Journal home page: <http://www.aensiweb.com/AEB/>

Feasibility Study of Urban Raw Wastewater for Agricultural Uses (Case Study: Meshkinshahr, Iran)

MahmoodNoori Mashiran and Ebrahim Fataei

Department of Environmental science, Ardabil Branch, Islamic Azad University, Ardabil, Iran

ARTICLE INFO

Article history:

Received 12 October 2014

Received in revised form 26 December 2014

Accepted 1 January 2015

Available online 17 February 2015

Key words:

quality of wastewater, raw urban wastewater, agricultural application, heavy metals, Meshkinshahr, Iran

ABSTRACT

Today, scarcity of water is one of the greatest challenges in agriculture. The reasons for this scarcity are population growth, promotion of lifestyle, climate change, continuous droughts, and lack of proper water resource management. Under such circumstances, use of unconventional water sources including urban wastewaters in agriculture as the major consumer of water is of special importance. In this paper, quality of raw wastewater of Meshkinshahr, located in Ardabil Province, Iran, was studied, which is directly used in spring and summer in Brazil Village, a village in the southern part of the county, due to shortages of water sources for irrigation of orchards and farms. The present research is a descriptive-cross sectional study carried out by sampling from raw urban wastewaters in spring and summer 2014. The results suggested that average values of pH, DO, COD, BOD₅, TSS, Cr, Mg, B, Cd, Pb, Hg, Fe, parasite eggs, oil and fat, turbidity, nitrates, Ca, Mn, Na, Cl and phosphates were 7.05 mg/l, 2 mg/l, 412.5 mg/l, 235 mg/l, 92.5 mg/l, 0.0265 mg/l, 0.02 mg/l, 0.15 mg/l, 0.015 mg/l, 0.22 mg/l, 0.05 mg/l, 0.275 mg/l, 4.8 mg/l, 91.5 mg/l, 0.085 mg/l, 53.5 mg/l, 37 mg/l, 39.5 mg/l, 0 mg/l, 12.55mg/l, respectively, which are lower than agricultural standards of Department of the Environment (DOE). Average contents total and fecal coliforms were 3300mg per100ml and 2300mg per100ml, respectively, which were inconsistent with this standard. Thus, in order to avoid hygienic and coliform-based impacts, chlorinating the wastewater prior to its consumption for agricultural purposes is recommended to reduce the values to DOE standards.

© 2015 AENSI Publisher All rights reserved.

To Cite This Article: MahmoodNoori Mashiran and Ebrahim Fataei., Feasibility Study of Urban Raw Wastewater for Agricultural Uses (Case Study: Meshkinshahr, Iran). *Adv. Environ. Biol.*, 9(2), 695-699, 2015

INTRODUCTION

Today, world is facing the challenge of water scarcity. Limitation of water sources has attracted attention of researchers to proper use of unconventional water sources such as brine water and urban and industrial effluents. Due to urban development and increase in water consumption, a large volume of water is produced from raw wastewater and treated effluents. In Iran, a great deal of water used by major cities is converted to urban wastewaters [1]. While many arid and semiarid areas of Iran are exposed to scarcity of water sources, and water demands for irrigation are high. Under these conditions, usage of low quality sources is being considered. On the other hand, the urbanized areas produced huge amounts of wastewater, the inappropriate disposal of which poses environmental problems to the surrounding areas. Application of urban wastewaters in irrigation has been recommended as a rich source of fertilizing elements in many countries [2].

Wastewater can compromise human health through incidence of bacteriological, viral, protozoan and parasitic diseases. Fatta *et al* [3] believed that application of wastewater in agriculture, provided it is sufficiently treated, will not cause nematode diseases among farm workers and consumers, unless climatic conditions and irrigation type prepares the environment for survival of nematode eggs. But, according to Shuval *et al* [4], endemic ascariasis and trichuriasis in the world is coincided with application of wastewater for irrigation. In their epidemiological studies, these researchers also concluded that the risk of outbreak of ascaris worms in children and adults, who used wastewater-irrigated crops, is very high. They also point to diseases such as cholera, typhoid and shigellosis as a result of wastewater usage in farming.

Corresponding Author: Ebrahim Fataei, Department of Environmental science, Ardabil Branch, Islamic Azad University, Ardabil, Iran.
E-mail: Ebfataei@gmail.com

In addition to pathogenic microorganisms, domestic and industrial wastewaters contain a variety of chemical compounds including medicines, hormones, antibiotics, hormone-affecting compounds, etc., which their long-term impacts particularly via application of effluents in agriculture on human health and ecosystems are yet to be known [3].

Generally, wastewater leaves various biological, physical and chemical impacts. These impacts mainly result from physico-chemical quality of wastewater, which in casefarms are irrigated with it, influence physical conditions of the soil of cultivated lands and assimilation capacity of micro and macro elements (Shariati, 1996). Irrigation with wastewater is taken into consideration as an option to offset shortage of available water. In this case, it is necessary to assess impacts of use of urban wastewater on physical and chemical properties to achieve sustainable development [6]. In Iran, based on a report by Shayegan and Afshari, out of 3.9 billion m³ urban wastewater, only 9% are treated and the rest enter absorption wells, rivers and farms without any treatment [7].

During last century, water consumption rate increased heavily. Increase in consumption per capita and unplanned use of water sources has led to qualitatively and quantitatively critical conditions of water. Hence, usage of unconventional water sources including urban wastewaters, particularly in agriculture, which is the main consumer of water, is of special importance. Many developing countries are not capable to use detailed wastewater treatment plans. Millions of farmers in these regions do the farming by using wastewater or wastewater-polluted sources, where there is no alternative for wastewater. Deleterious components in the wastewater can threaten environmental health and quality. Risk management and intermediate solutions are, therefore, necessary to prevent from adverse impacts in wastewater irrigation. A combination of control measures including control at source, control at farm level, and postharvest measures are applied to protect farmers and consumer [7].

Due to high potential to hurt human and animal health, heavy elements such as Pb, Cd and Cr were highly taken into consideration in recent years and efforts have been made to prevent them from entering environmental cycles as much as possible [8].

Since 1960, some studies were carried out about health hazards of projects using wastewater in irrigation. Also in 1968, wide research was conducted in Namibia with the aim of preparing drinking water from wastewater. Then, in 1970s and 1980s, potential of health hazards in reusing treated wastewater for irrigation and drinking purposes. In the last quarter of 20th century advantages of wastewater reuse was officially confirmed as an option in water resource development and became legitimate by United States and European Union. Use of wastewater for irrigation (due to high concentrations of heavy and toxic metals and their microbiological load) can disrupt ecosystem functioning [9].

Many developing countries are not able to use detailed treatment plans. Millions of farmers in the suburban areas of countries do the farming by using wastewater or waters polluted with agricultural wastewaters and they have no alternative. Deleterious components of wastewater can threaten environmental health and quality. Risk management and intermediate solutions are necessary to prevent adverse impacts of wastewater. A combination of control measures including control at source, control at farm level, and postharvest measures to protect farmers and consumers [10].

In "study of the impacts of irrigation with urban wastewater on soil and growth of pine trees in Tehran" Salehi *et al* [12] concluded that utilization of urban wastewater could be a source of water for afforestation.

Naseri [11] in "study of methods, health criteria, and wastewater reuse project management" conclude that considerable amounts of materials such as phosphate, potassium hydroxide, and Nitrogen, which all play a vital role in fertilizing arable lands, effectively increase crop production. On the other hand, new lands will be cultivated in order to supply water for agriculture leading to emigration of the farmers.

MATERIALS AND METHODS

According to existing meteorological data, average annual rainfall in Meshkinshahr City located west of Ardabil Province, northwest of Iran is 390mm. This city lies in a mountainous area with temperate and relatively cold weather and most crops are irrigated, which is exclusively supplied by surface waters, wells, and springs.

This is descriptive-cross sectional study that sampling from raw urban wastewater was directly performed in orchards and farms in spring and summer to determine levels of important parameters in irrigating agricultural crops in orchard and farm inlets in spring and summer (May to August). Sampling was carried out based on Standard Method. Samples were analyzed in creditable laboratories of Ardabil Province (Okinchi) and East Azerbaijan (Haghdar in Miyaneh and Kolor Pars). Samples for microbial and biological tests were kept in 4° C and were transferred to the laboratory in at most 6 hours. The samples for heavy metal measurement were transferred to the laboratory after nitric acid was added. Some parameters including DO, temperature, and pH were measured onsite by portable 156 sension hach. Excel and SPSS16 were used to draw the graphs and statistical analyses.

RESULTS AND DISCUSSION

A variety of standards about different applications of effluent has been set by international organizations such as FAO, EPA, and WHO. In Iran, the standards of the use of wastewater in agriculture and irrigation were set by DOE. Decisions on potentials of wastewater reuse I various options have been made based on results of

tests conducted on raw urban wastewaters in Meshginshahr and its comparison to DOE standards (Table1). Since the values obtained in four times of sampling were not considerably different, the average measured values (Table1) were compared to national standards.

Table 1: A comparison of the quality of raw urban wastewater in Meshginshahr with DOE standards for irrigation and agricultural purposes.

DOE standards for irrigation and agricultural purposes	average measured values	unit	Parameter	No
6-8.5	7.05	-	pH	1
2	2	mg/L	Do	2
100	235	mg/L	BOD ₅	3
200	412.5	mg/L	COD	4
100	92.5	mg/L	TSS	5
-	619	s/cm μ	EC	6
-	420.5	mg/L	TDS	7
-	1.0126	ML/LIT	SAR	8
0.1	0	mg/L	Remaining chlorine	9
400	2300	MPN/100mL	Fecal coliform	10
1000	3300	MPN/100mL	Total coliform	11
Bylaws 5	0	Count	Parasit eggs	12
-	53.5	mg/L	Ca	13
100	37	mg/L	Mg	14
-	39.5	mg/L	Na	15
-	11.8	mg/L	Nitrate	16
-	12.55	mg/L	Phosphate	17
1	0.0265	mg/L	Cr	18
1	0.15	mg/L	Br	19
1	0.02	mg/L	Mn	20
0/05	0.015	mg/L	Cd	21
1	0.22	mg/L	Pb	22
little	0.05	mg/L	Hg	23
50	91.5	NTU	Turbidity	24
10	4.8	mg/L	Fat and oil	25
1	0	mg/L	Phenol	26
-	52.95	mg/L	Ammonia	27
3	0.275	mg/L	Fe	28
-	0.085	mg/L	Nitrite	29
4	11	mg/L	Chloride	30

As seen in Table1, average values of total coliform and fecal coliform in urban wastewaters in Meshginshahr were 3300mg/l and 2300 mg/l, respectively. Therefore, these values are far beyond the values recommended by DOE (1000mg/l, 400mg/l). Average values of pH, DO, COD, BOD₅, TSS, Cr, Mg, B, Cd, Pb, Hg, Fe, parasite eggs, oil and fat, turbidity, nitrates, Ca, Mn, Na, Cl and phosphates were 7.05 mg/l, 2 mg/l, 412.5 mg/l, 235 mg/l, 92.5 mg/l, 0.0265 mg/l, 0.02 mg/l, 0.15 mg/l, 0.015 mg/l, 0.22 mg/l, 0.05 mg/l, 0.275 mg/l, 4.8 mg/l, 91.5 mg/l, 0.085 mg/l, 53.5 mg/l, 37 mg/l, 39.5 mg/l, 0 mg/l, 12.55mg/l, respectively, which are consistent with DOE standards. Results of sampling for these parameters in comparison with DOE standards are grouped separately as heavy metals (Diag.1), biological parameters (Diag.2), nutrients (Diag.3) and anions and cations (Diag.4).




Diagram 1: Comparison of heavy metal (Cd, Pb, Hg, Cr, Mn, and B) contents in Meshkinshahr with DOE standards for usage of wastewater in agriculture.

Selection of suitable plants to be irrigated by wastewater should follow some principles to avoid infection of the crops with pathogens and transferring them to consumers. Thus, considering direct use of raw wastewater in Meshkinshahr and the obtained information of its quality, irrigating vegetables and crops, which are eaten uncooked, is not recommended. Also due to direct contact with soil, irrigating rooted plants such as potato, wheat, barley and beans, which are consumed directly, is not recommended. Therefore, when raw wastewater is conveyed to the treatment plant and coliform contents are reduced to a standard level for agricultural purposes, effluents of Meshkinshahr treatment plant could be used in agriculture.




Diagram 2: Comparison of BOD₅, COD and TSS with DOE standards in Meshkinshahr wastewater.




Diagram 3: Contents of Nutrients (phosphate and nitrate) in Meshkinshahr wastewater.




Diagram 4: Contents of Ca²⁺, Mg²⁺ and Na⁺ in Meshkinshahr wastewater.

Irrigation of orchards, where fruits are consumed freshly; trees, which their crops are used in the form of nuts; wooden trees including cedars, pines, elms, and spruces, as well as industrial crops such as cotton are not recommended because of soil contamination. Irrigating green fields and public places are not recommended either. Although concentrations of heavy metals are lower than the thresholds recommended by DOE, accumulative impacts of these elements make irrigating nonedible industrial crops such as cotton and wooden trees. At present, direct and unplanned use of raw wastewater in downstream areas of Meshginsahr is for farming, hence, in order to protect farmers' and villagers' health, these sources should be available for farmers in a planned manner by transmission to the treatment plant to be chlorinated to meet DOE standards regarding total and fecal coliforms.

Conclusions:

The results showed that average values of pH, DO, COD, BOD₅, TSS, Cr, Mg, B, Cd, Pb, Hg, Fe, parasite eggs, oil and fat, turbidity, nitrates, Ca, Mn, Na, Cl and phosphates were 7.05 mg/l, 2 mg/l, 412.5 mg/l, 235 mg/l, 92.5 mg/l, 0.0265 mg/l, 0.02 mg/l, 0.15 mg/l, 0.015 mg/l, 0.22 mg/l, 0.05 mg/l, 0.275 mg/l, 4.8 mg/l, 91.5 mg/l, 0.085 mg/l, 53.5 mg/l, 37 mg/l, 39.5 mg/l, 0 mg/l, 12.55mg/l, respectively, which are consistent with DOE standards for use of wastewater in irrigation and agriculture. Only average contents of total and fecal coliforms were 3300mg per 100ml and 2300mg per 100ml, which was higher than DOE standards.

REFERENCES

- [1] Tajrishi. M., 1998. A new and comprehensive approach to wastewater challenge in Tehran City, Water and wastewater, 28: 30-37.
- [2] Feigin, A., I. Ravina and J. Shalheret, 1991. Irrigation with treated sewage effluent. Berlin:Springer-Verlag.
- [3] Fatta, D. and N. Kythreotou, 2005. Wastewater as Valuable Water Resource - Concerns, Constraints and Requirements Related to Reclamation, Recycling and Reuse. IWA International Conference on WaterEconomics, Statistics, and Finance, Greece.

- [4] Shuval, H.I., A. Adin, B. Fattal, E. Rawitz, P. Yekutieli, 1986. Wastewater Irrigation in Developing Countries. Health Effects and Technical Solutions. World Bank Tech. Pap, 51: 325.
- [5] Shariati, M.R., 1996. Investigation of chemical quality and use of wastewater in irrigation. Journal of Water, Soil and Environment, 10: 51-55.
- [6] MasoudiAshtiyani, S., M. Parsinezhad and F. Abbassi, 2011. Effects of urban irrigating *sorghum* with wastewater on physical properties of soil. Journal of Soil research (Soil and water sciences), 25(3): 25-33.
- [7] Shayegan, J. and A. Afshari, 2004. Study of urban and industrial wastewaters of Iran, Journal of Water and wastewater, 49: 58-69.
- [8] Sahebghadam, A., 1988. Lead metabolism and its resulting poisoning. TarbiatModaress Press.
- [9] Naghshinepour, B., 1994. Application of wastewater in agricultural production and soil adjustment. First congress on infrastructural (water and soil) planning and policy making, Ministry of Agriculture, Tehran: 135-144.
- [10] Qadir, M., D. Wichelns, L. Raschid-Sally, P.G. McCornick, P. Drechsel, P. Drechsel, A. Bahri, P.S. Minhas, 2010. The challenges of wastewater irrigation in developing countries. Agricultural WaterManagement, 97(4): 561-568.
- [11] Naseri, A.A., S.A. MohseniMovahed, H.A. Kashkooli, Zare' Abyaneh and M.R. Ehasani, 1999. Assessment of empirical methods of estimating evapotranspiration potentials in Hamadan. Agricultural Researches, Scientific-research Journal of faculty of agriculture, Abu Ali Sina University.
- [12] SalehiArjmand, H., M.H. Mahdyan, A.R. Kargari, M. Mahdye, 2007. Feasibility study of using effluents of Arak treatment plant in agriculture. Water and environment, 65: 41-46.