

Effect Of Carbon Sources On *Bacillus sp.R2* Chitinase Production

^{1,2,3}Ben Amar Cheba, ³Taha Ibrahim Zaghloul, ⁴Ahmad Rafik EL-Mahdy, ³Mohamad Hisham EL-Massry

¹Department of Biology, College of Science, Al Jouf University, Kingdom of Saudi Arabia (KSA).

²Department of Biotechnology, Faculty of Nature and Life Sciences, University of Sciences and Technology of Oran -Mohamed Boudiaf (USTOMB), BP 1505 Al Mnaouar, Oran 31000, Algeria.

³Department of Biotechnology, Institute of Post Graduate Studies and Research, University of Alexandria- Alexandria –Egypt.

⁴Department of Food Science and Technology-Faculty of Agriculture-University of Alexandria- Alexandria-Egypt.

Address For Correspondence:

Ben Amar Cheba ,Department of Biology, College of Science, Al Jouf University, Kingdom of Saudi Arabia (KSA).
Phone number 0966 536832385; E-mail: benach57@yahoo.com

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Received 8 January 2017; Accepted 28 April 2017; Available online 24 May 2017

ABSTRACT

Effect of carbon sources on *Bacillus sp.r2* chitinase production were determined using one variable at time technique (OVAT) in shake flasks (submerged fermentation). The tested factors were chitin sources and forms, colloidal chitin concentration, secondary carbon sources and glucose concentration. The highest chitinase activity (57 U/ml) was obtained in 75% natural sea water, 0.5% shrimp shell colloidal chitin as best and essential carbon source, 0.5% glucose. The production also was optimum medium volume 50 ml/250ml flask and 24 h incubation period at 180 rpm shaking. carbon sources affect strangely enzyme induction and may help to understand the physiology of enzyme production

KEYWORDS: *Bacillus sp. R2*, Carbon Sources chitinase, production, OVAT, optimization.

INTRODUCTION

Chitin, a β - [1-4] homopolymer of N-acetyl - D- glucosamine (Glc NAc), is the second most abundant polysaccharide existing in nature after cellulose. It is a major structural component of most biological systems such as mollusks, insects, crustaceans, fungi, algae and marine invertebrates [1]. Chitin and its derivatives are of commercial and biotechnological interest because they have various biological activities and wide range of applications in areas ranging from waste water treatment to agrochemical and biomedical uses [2,3,4,5]. Chitinases (E.C.3.2.1.14), which are found in variety of organisms such as viruses, bacteria, actinomycetes, yeasts, fungi, plants, animals and also in human beings[6,7]. During the last decade, chitinases have received an increased attention due to their wider ranges of biotechnological applications especially in the biocontrol of fungal phytopathogens[8] and harmful insects[9]. They have also been used as vaccine [10], antitumor[11], tumor marker[12] and useful biomarker in Gaucher disease[13] and used in the preparation of sphaeroplasts and protoplasts from yeasts and fungal species which can be used further for biotechnological uses[14]. In the previous studies, *Bacillus sp.R2* was screened as hyper chitinase producer and identified. In the present work the effect of carbon sources on *Bacillus sp.r2* chitinase production will be determined using one variable at time technique (OVAT) (OVAT) in shake flasks.

MATERIAL AND METHODS

Substrates and chemicals:

Chitin was extracted from crustaceans by the method of [15], Crab shell chitin flakes (Win-lab, UK). Swollen chitin was prepared according Monreal and Reese, [16]. Peptone tryptone, and yeast extract were obtained from (Oxoid Hampshire, England). 2 Hydroxy 3,5 dinitrosalselic acid (DNSA) obtained from (Merck, Darmstadt- Germany);. N-acetyl glucosamine, agarose, bovine serum albumin (BSA) were from (Sigma -USA). All other chemicals used were of the highest grade available.

Bacterial strain and culture condition:

Bacillus sp.R2 was newly marine bacterial strain isolated from red sea Egypt, identified either by biochemical tests or 16S rRNA methodology (strain accession number in NCBI GenBank was: DQ923161). To maintain the isolated bacterial cells the short term maintenance was performed repeatedly at an interval of 2-3 months at 4°C using marine LB Agar slants. Moreover, the long term maintenance, more than 2 years, was performed by adding 0.5 ml of the early stationary phase cultures grown in marine LB to 50% (v/v) sterile glycerol and the cultures were kept at -20°C. For chitinase production several single colonies of the strain Bacillus sp. R2 were used to inoculate 5 ml marine LB medium supplemented with 0.5% colloidal chitin. This bacterial culture was allowed to grow at 37°C for 24 h using orbital shaking incubator, Overnight seed culture 2,5% (v/v) was used to inoculate 50 ml production medium The culture was incubated at 30°C for 24 h at 180 rpm shaking.

Optimization of chitinase production:

Effect of carbon sources on Bacillus sp.r2 chitinase production were determined using one variable at time technique (OVAT) in shake flasks (submerged fermentation) .The tested factors were chitin sources and forms, colloidal chitin concentration, secondary carbon sources and glucose concentration.

Analytical procedures:

Growth monitoring and Protein assay:

Colony forming units (CFU) was determined [17]. Moreover, bacterial growth was monitored spectrophotometrically by measuring the absorbance of the cultures at 660 nm. Soluble proteins were determined as described by Bradford [18] using bovine serum albumin as standard

Chitinase assay:

Chitinase activity was analyzed by estimating the released reducing ends of sugar according to the method of Miller [19] using N-acetyl - D-glucosamine (NAG) as standard .One unit of chitinase activity was defined as the amount of enzyme required to release 1 μ mol of NAG per minute during these conditions.

RESULTS AND DISCUSSION

Optimization of chitinase production:

Effect of chitin sources and forms:

Chitinase production was remarkably affected by the nature and form of chitin. The shrimp shell colloidal chitin was the best substrate followed by Sigma crab shell chitosan and prawn shell colloidal chitin powder (Table1). Chitin flasks were not as effective as colloidal chitin. This may be due to the open structure of colloidal chitin which makes it more accessible by the enzyme. Results were in agreement with those reported about chitinase inducibility by colloidal chitin [20,21,22,23]. These chitins still vary however in strand opening (degree of compactness), degree of acetylation and probably the presence of covalently linked components other than N-acetyl glucosamine.

Effect of colloidal chitin concentration:

It is known that an ideal substrate concentration in any fermentation process results in a higher conversion efficiency and an optimum substrate utilization [24]. Based on above results Figure (1) colloidal chitin was the most suitable substrate for chitinase production. Various concentrations of colloidal chitin was tested in order to obtain the maximum chitinase production, the optimum concentration for chitinase production was 0.5%. Many investigators reported that the optimal concentration ranged between 0.5 and 1 % [25,22,26,27].

Effect of carbon sources:

Various carbon source supplementations were tested to investigate their effect on the chitinase production. Glucose, N-acetylglucosamine, maltose and pectin showed significant increase in chitinase production based on units/mg protein, respectively (Table 2). On the other hand, glycerol, Tween 20 and Tween 80 gave only maximum growth rate and protein content. Other carbon sources did not promote the enzyme production. Many

workers found that, the addition of glucose, glucosamine or N-acetylglucosamine, stimulated chitinase production. [28,29,30,22,31,32] Moreover, others reported that, the addition of maltose [33], β -cyclodextrin [34], pectin [35] or CMC [36] also enhanced significantly the chitinase production.

Effect of glucose concentration:

Since the addition of glucose stimulates the chitinase production, attempts were made to determine the suitable glucose concentration. The best concentration of glucose was found to be 0.5% which gave maximum induction level after 24 hrs incubation. However, concentrations more than 0.5% showed an adverse effect (Figure 2). It was reported that the optimal glucose concentration for chitinase production ranged between 0.2 and 1%. It was 0.2% for *Acinetobacter sp* [30], 0.3% for *Aspergillus sp.* [25], 0.5% for *Bacillus sp.* BG11 [31] and *B. licheniformis*[22], 0.6% for *Vibrio alginolyticus*[28] and 1% for *Salinivibrio costicola*(29). Furthermore, it was noticed for marine bacteria that glucose, trehalose and chitin hydrolysis end products e.g., N-acetyl glucosamine, glucosamine and chitobiose play an important role in chemotaxis and induce the chitinase production [37,38,39].

Table 1: Effect of chitin sources and forms on chitinase production by strain R2.

Substrate sources	Substrate forms	Protein (mg/ml)	Activity (U/ml)	Specific Activity (U/mg)
Crab shell chitin-flasks (Win lab.UK)	α	0.095	9.66	101.4
Crab shell chitin-powder (Winlab.UK)	α	0.105	11.4	108
Crab shell chitin	α	0.081	9.32	115
Shrimp shell chitin	α	0.060	8	131.9
Prawn shell chitin	α	0.071	10.3	143.8
Squid chitin	β	0.075	14.2	189.3
Crab shell colloidal chitin (Win lab)	α	0.125	24.1	191.5
Crab shell swollen chitin	α	0.129	21.2	163.7
Crab shell colloidal chitin	α	0.118	20.6	174.4
Shrimp shell colloidal chitin	α	0.133	31.3	234.1
Prawn shell colloidal chitin	α	0.153	32	208.2
Crab shell chitosan (Sigma - USA)	α	0.103	22.7	219.5
Crab shell chitosan	α	0.106	19.5	182.6
Crab shell colloidal chitosan	α	0.131	13	98.5




Fig. 1: Effect of substrate concentration on chitinase production

Table 2: Effect of carbon source on chitinase production

Carbon source	Conc. (%)	Growth OD _{660nm}	Chitinase activity (U/ml)	Protein (mg/ml)	Specific activity (U/mg)
Monosaccharides					
Glucose	0.5	1.94	48.02	0.115	417
Galactose	0.5	1.87	14.61	0.143	102.16
Fructose	0.5	1.94	5.25	0.117	44.87
Sorbitose	0.5	1.52	12.86	0.129	99.6
Arabinose	0.5	1.93	21.21	0.132	160.68
Xylose	0.5	1.92	2.42	0.112	21.60
Mannose	0.5	1.68	0.53	0.096	5.52
Disaccharides					
Maltose	0.5	1.89	37.91	0.122	310.77
Lactose	0.5	1.81	12.45	0.091	136.81
Sucrose	0.5	1.78	7	0.089	78.65
Aminosugar(NAc glucosamine)	0.5	1.93	41.41	0.108	383.42
Polyols (glycerol)	0.5	2.23	6.12	0.105	58.28

Glucose based polysaccharides					
Starch	0.2	1.91	36.76	0.124	296
Dextran	0.2	1.75	0.134	0.071	1.88
CMC	0.2	1.64	2.02	0.062	32.58
Cellulose	0.2	1.52	1.75	0.053	33.01
Galactose based polysaccharides					
Agar	0.1	1.73	6.80	0.078	87.17
Agarose	0.1	1.85	10.23	0.081	126.29
Pectin	0.2	1.87	30.43	0.104	292.59
Arabic gum	0.2	1.61	8.48	0.075	113.06
Miscellaneous					
Alginate	0.2	1.57	13.67	0.087	157.12
Tween 20	0.2	2.36	7.81	0.238	32.81
Tween 80	0.2	2.58	10.16	0.250	40.64




Fig. 2: Effect of glucose concentration on chitinase production

Conclusion:

The optimization of medium composition specially carbon sources which affect strangely enzyme induction and plays a crucial role in the microbial production of chitinases and help to understand the physiology of enzyme production, recognizing the key factors, setting the best bioprocess (fermentation) conditions which pave the way for the industrial scale-up with low cost and high enzymatic yield.

ACKNOWLEDGEMENT

Thanks are due to the Algerian ministry of higher education and scientific research for their financial support. also the author thanks Professor Almaahdy R.A for kindly providing some chemicals and his useful scientific discussion. We are grateful to Dr. Zaghoul T.I. and Dr. EL-Massry M.H for their scientific assistance and helpful suggestions and comments during the work.

REFERENCES

- [1] Zargar, V., M. Asghari and A. Dashti, 2015. A Review on Chitin and Chitosan Polymers: Structure, Chemistry, Solubility, Derivatives, and Applications. *Chem.Bio.Eng. Reviews*, 2: 204-226.
- [2] Van den Broek, L.A.M., R.J.I. Knoop, F.H.J. Kappen, C.G. Boeriu, 2015 Chitosan films and blends for packaging material. *Carbohydr. Polym.*, 116: 237-242.
- [3] Guilherme, M.R., F.A. Aouada, A.R. Fajardo, A.F. Martins, A.T. Paulino, M.F.T. Davi, A.F. Rubira, E.C. Muniz, 2015. Superabsorbent hydrogels based on polysaccharides for application in agriculture as soil conditioner and nutrient carrier: A review. *European Polymer Journal*, 72: 365.
- [4] Usman, A., K.M. Zia, M. Zuber, S. Tabasum, S. Rehman, F. Zia, 2016 Chitin and chitosan based polyurethanes: A review of recent advances and prospective biomedical applications. *Int J Biol Macromol.*, 86: 630-45.
- [5] Rafique, A., K. Mahmood Zia, M. Zuber, S. Tabasum, S. Rehman, 2016. Chitosan functionalized poly(vinyl alcohol) for prospects biomedical and industrial applications: A review. *Int J Biol Macromol.*, 87: 141-54.
- [6] Di Rosa, M., G. Distefano, K. Zorena, L. Malaguarnera, 2016 Chitinases and immunity: Ancestral molecules with new functions. *Immunobiology*, 221(3): 399-411.

- [7] Kzhyshkowska, J., S. Yin, T. Liu, V. Riabov, I. Mitrofanova, 2016. Role of chitinase-like proteins in cancer. *Biol Chem.*, 397(3): 231-47.
- [8] Nagpure, A., B. Choudhary, R.K. Gupta, 2014. Chitinases: in agriculture and human healthcare. *Crit Rev Biotechnol.*, 34(3): 215-32.
- [9] Weixing Zhu, Di Wang, Tian Liu and Qing Yang, 2016. Production of N-Acetyl-d-glucosamine from Mycelial Waste by a Combination of Bacterial Chitinases and an Insect N-Acetyl-d-glucosaminidase J. *Agric. Food Chem.*, 64(35): 6738-6744.
- [10] Elieh-Ali-Komi, D., M.R. Hamblin, 2016. Chitin and Chitosan: Production and Application of Versatile Biomedical Nanomaterials. *Int J Adv Res (Indore)*; 4(3): 411-427.
- [11] AV Il'ina, VP Varlamov, 2015. In vitro antitumor activity of heterochitooligosaccharides (Review). *Applied biochemistry and microbiology*, 51(1): 5-14.
- [12] Hamilton, Gerhard, Barbara Rath and Otto Burghuber, 2015. "Chitinase-3-like-1/YKL-40 as Marker of Circulating Tumor Cells." *Translational Lung Cancer Research*, 4(3): 287-291.
- [13] Cheba, B., T.I. Zaghoul, A.R. EL-Mahdy and M.H. EL-Massry, 2011. Enhanced production of *Bacillus sp.* R2 chitinase through cell immobilization. *ACT-Biotechnology Research Communications*, 1(1): 8-13.
- [14] Nagpure, A., B. Choudhary, R.K. Gupta, 2014. Chitinases: in agriculture and human healthcare-reviews in biotechnology, 34(3): 215-232.
- [15] Synowiecki, J., Z. Sikorski and K.M. Nacz, 1982. Immobilisation of amylases on krill chitin. *Food chem.*, 8: 239-246.
- [16] Monreal, J and E.T. Reese, 1969. The chitinase of *Serratia marcescens*. *Can. J. Microbiol.*, 15: 689-696.
- [17] Pelczar, M.J. and E.C. Chan, 1977. *Laboratory exercises in microbiology*, 4th edition, Mc Graw Hill, nc
- [18] Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- [19] Miller, G.R., 1959. Use of Dinitrosalicylic Acid reagent for determination of reducing sugar. *Anal. Chem.*, 31(3): 426-428.
- [20] Samac, D.A., C.M. Hironaka, P.E. Yallaly and D.M. Shah, 1990. Isolation and characterization of genes encoding basic and acidic chitinases in *Arabidopsis thaliana*. *Plant Physiol.*, 93: 907-914.
- [21] Watanabe, T., W. Oyanagi, K. Suzuki and H. Tanaka, 1990. Chitinase system of *Bacillus circulans* WL-12 and importance of chitinase Al in chitin degradation. *J. Bacteriol.*, 172 (7): 4017-4022.
- [22] Ohishi, K., M. Yamagishi, T. Ohta, M. Suziki, H. Izumida, H. Sano, et al., 1996. Purification and properties of two chitinases from *Vibrio alginolyticus* H-8. *J. Ferment. Bioeng.*, 82: 598-600.
- [23] El-sayed, E., S.M. Ezzat, M.F. Ghaly, M. Mansour and M.A. El-Bohey, 1999. Purification and characterization of two chitinases from *Streptomyces albobovineus* S-22. *J. Union Arb. Biol.* 8 (B). *Microbiology and viruses* 149-163, 6th international conference, 8-11.
- [24] Montgomery, M.T., and D.L. Kirchman, 1994. Induction of chitin-binding proteins during the specific attachment of the marine bacterium *Vibrio harveyi* to chitin. *Appl. Environ. Microbiol.*, 60: 4284-4288.
- [25] Rattanakit, N., A. Plikomol, S. Yano, M. Wakayama and T. Tachiki, 2002. Utilization of shrimp shell fish waste as a substrate for solid-state cultivation of *Aspergillus sp.* S1-13: Evaluation of a culture based on chitinase formation which is necessary for chitin assimilation. *J. Biosc. Bioeng.*, 93(6): 550-556.
- [26] Wen, C., C. Teseng, C. Cheng and Y. Li, 2002. Purification, characterization and cloning of a chitinase from *Bacillus sp.* NTU2. *Biotechnol. App. Biochem.*, 35: 213-219.
- [27] Yuli, P.E., M.T. Suhartono, Y. Rukayadi, J.K. Hwanj and Y.R. Pyun, 2004. Characteristic of thermostable chitinase enzymes from the Indonesian *Bacillus sp.* 13.26. *Enz. Microb. Technol.*, 35: 147-153.
- [28] Aunpad, R., and W. Panbamgrad, 2003. Cloning and characterization of the constitutively expressed chitinase C gene from a marine bacterium *Salinivibrio costicola* strain 5SM-1. *J. Biosc. Bioeng.*, 96: 529-536.
- [29] Cabib, E., 1988. Assay for chitinase using tritiated chitin in: Wood WA, kelloggst, editors. *Methods in enzymol.* Vol 61 San Diego CA: Academic press, p: 424-426.
- [30] Shimosaka, M., M. Nogawa, X. Wang, M. Kumehara and M. Okazaki, 1995. Production of two chitosanases from a chitosan-assimilating bacterium, *Acinetobacter sp.* strain CHB101. *App. Environ. Microbiol.*, 61(2): 438-442.
- [31] Sakai, K., A. Yokota, H. Kurokawa, M. Wakayama and M. Moriguchi, 1998. Purification and characterization of three thermostable endochitinases of a noble *Bacillus* strain, MH-1 isolated from chitin-containing compost. *App. Environ. Microbiol.*, 64(9): 3397-3402.
- [32] Rattanakit, N., A. Plikomol, S. Yano, M. Wakayama and T. Tachiki, 2002. Utilization of shrimp shell fish waste as a substrate for solid-state cultivation of *Aspergillus sp.* S1-13: Evaluation of a culture based on chitinase formation which is necessary for chitin assimilation. *J. Biosc. Bioeng.*, 93(6): 550-556.
- [33] Liu, B.L., P.M. Kao, Y.M. Tzeng and K.C. Feng, 2003. Production of chitinase from *Verticillium lecanii*. *Enz. Microb. Technol.*, 33: 110-115.
- [34] Shimosaka, M., M. Nogawa, X. Wang, M. Kumehara and M. Okazaki, 1995. Production of two

- chitosanases from a chitosan-assimilating bacterium, *Acinetobacter* sp. strain CHB101. *App. Environ. Microbiol.*, 61(2): 438-442.
- [35] Chen, H.C., M.F. Hsu and S.T. Jiang, 1997. Purification and characterization of an exo-N, N-diacetylchitobiohydrolase - like enzyme from *Cellulomonas flavigena* NTOU1. *Enz.Microb. Technol.*, 20: 191-197.
- [36] Melchers, L.S., M. Apotheker De-Groot, J.A. Van-Der knaap, A.S. Ponstein, Sela. M.B. Burlage, J.F. Bol, B.J.C. Cornelissen, P.J.M. Van-Der Elzen and H.J.M. Linthost, 1994. A new class of tobacco chitinases homologous to bacterial exo-chitinases displays antifungal activity. *Plant., J.* 5: 569-580.
- [37] Sheng, L., Z. Zhi-an, Ming Li., G.V. Zhen-Rong, B. Chen and D.H. Wei, 2002. Purification and characterization of a novel chitinase from *Bacillus brevis*. *Acta. Biochem. Biophys. Sinica B4*, (6): 690-696.
- [38] Yu, C., B.L. Bassler and S. Roseman, 1993. Chemotaxis of the marine bacterium *Vibrio furnissii* to sugars: A potential mechanism for initiating the chitin catabolic cascade. *J. Biol. Chem.*, 288(13): 9405-9409.
- [39] Yu, C., A.M. Lee and S. Roseman, 1987. The sugar specific adhesion/deadhesion apparatus of the marine bacterium *Vibrio furnissii* is a sensorium that continuously monitors nutrient levels in the environment. *Biochem. Biophys. Res .Commun.*, 149: 86-92.